Standard Immunisation catch-up program template For children currently aged from 2 years up to 4 years (<4 years) with <u>NO</u> documented history of vaccination

Important information

- There is the option of using the '*Accelerated* Immunisation Catch-up Template for children aged from 2 years up to 4 years' [<4 years]
- To prevent Centrelink payment issues complete an Immunisation History Form. Fill out Part A, Part B [only need to ✓ 'on a catch-up program'] and sign Part C then fax to Australian Immunisation Register (AIR) Fax no: 08 9254 4810.
- If a child has a DOB on or after 01/10/2014 to be considered up to date they will need to catch up for the Infanrix or Tripacel [DTPa] booster dose scheduled at 18 months. In a catch-up program Infanrix or Tripacel always need to be a minimum of 6 months spacing from any other diphtheria/tetanus/pertussis [DTPa] containing vaccine such as Infanrix Hexa or Infanrix-IPV.
- Sources for checking vaccination records: AIR records; VIVAS records; Personal Health Record Book; patient records; previous service provider.

Due now:	Infanrix Hexa [DTP-hepB-IPV-Hib vaccine] dose 1 M-M-R II or Priorix [Measles/Mumps/Rubella vaccine] dose 2
Then, one month later give:	Infanrix Hexa [DTP-hepB-IPV-Hib vaccine] dose 2 Priorix Tetra or ProQuad [Measles/Mumps/Rubella/Varicella vaccine] dose 2
Then, one month later give:	Menitorix [Meningococcal C/Hib vaccine] dose 1
Then, two months later give:	Infanrix Hexa [DTP-hepB-IPV-Hib vaccine] dose 3 [give at least 3 months after the last dose of Infanrix Hexa]
Then, six months later give: [if is born from 01/10/201 <i>will need this first booster dose to be up to date on AIR</i>]	Infanrix or Tripacel [DTPa vaccine] dose 4 [give at least 6 months after dose 3 of Infanrix Hexa and always at least 6 months before the Infanrix-IPV].
Return at 4 years of age and give:	Infanrix IPV [DTP-IPV vaccine] dose 5 [if <u>did not</u> receive Infanrix or Tripacel label dose 4] [give at 4 years of age or least 6 months after the last dose of DTP containing vaccine e.g. Infanrix – Hexa, whichever is later]

Refer to the electronic version of the Australian Immunisation Handbook (AIH) 10th edition Table 2.1.7: Minimum acceptable dose intervals for children <10 years.

Contact the Sunshine Coast Public Health Unit for advice on the extra vaccines recommended for:

- Aboriginal and Torres Strait Islander children
- Premature [<32 weeks gestation] and/or low birth weight [<2000g] children
- Children identified as being at medical risk for invasive pneumococcal disease [IPD]. Refer to the Australian Immunisation Handbook 10th ed. [AIH] pps 326-327 for a list of conditions associated with an increased risk of IPD